

SPECIMEN LABEL

3336™ F

TURF AND ORNAMENTAL SYSTEMIC FUNGICIDE

For the prevention and control of turf diseases and the diseases of annual and perennial flowers, bedding plants, foliage plants, ground covers, plus deciduous and evergreen trees and shrubs.

ACTIVE INGREDIENT:

Thiophanate-methyl (dimethyl 4,4'-o-phenylenebis[3-thioallophanate]) 41.25%

INERT INGREDIENTS 58.75%

TOTAL 100.00%

This product contains 4.0 lb thiophanate-methyl per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

- IF SWALLOWED: Induce vomiting immediately by giving two glasses of water and sticking finger down throat. Contact a physician or poison control center. Never give anything by mouth to an unconscious person.
- IF IN EYES: Immediately flush eyes with plenty of water for at least 15 minutes. Get medical attention if irritation persists.
- IF ON SKIN: Wash exposed areas with soap and water. Seek medical attention as needed.
- IF INHALED: Remove exposed person to fresh air. If not breathing, give artificial respiration. Get medical attention.

PRECAUTIONARY STATEMENTS

● HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled, or absorbed through the skin. May cause irritation of eyes, nose, throat, and skin. Avoid contact with eyes and skin. Avoid breathing spray mist.

● PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical resistance selection chart.

WPS Uses: Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard, 40 CFR Part 170, must wear: long sleeve shirt and long pants, chemical resistant gloves (such as barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton ≥ 14 mils) and shoes with socks. For exposures in enclosed areas: a respirator with either an organic vapor-removing cartridge with a pre-filter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval prefix TC-14G) or a NIOSH approved respirator with an organic vapor (OV) cartridge or canister with any N, R, P, or HE prefilters.

NON-WPS Uses: Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard, 40 CFR Part 170, should wear: long-sleeved shirt and long pants, and shoes with socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

● ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by disposal of equipment washwaters. Do not apply, allow to drift, or drain or flush equipment onto non-target areas.

● STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in the original container in a dry, temperature controlled area. Do not store in a manner where cross-contamination with other pesticides, fertilizers, food or feed could occur. If spilled during storage or handling, contain/re-capture spillage and dispose of in accordance with the Pesticide Disposal Instructions listed below.

Pesticide Disposal: Wastes resulting from the use of this product are toxic and may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Do not re-use empty container. Triple rinse (or equivalent), then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.



CLEARY CHEMICAL CORPORATION
178 RIDGE ROAD
DAYTON, NJ 08810-1501

EMERGENCY PHONE NUMBERS:
M-F 9AM-5PM ET
800-524-1662 • 732-329-8399
24 Hour CHEMTREC
800-424-9300

Effective Date: 8/99
Replaces: 12/98

EPA Reg. No. 1001-69

10M3/00

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling, and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Exemption: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, Chemical resistant gloves (such as barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, polyvinyl chloride (PVC) \geq 14 mils, or Viton \geq 14 mils), and Shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box only apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Turf and Landscape Uses: Keep children and pets out of treated areas until sprays have dried.

GENERAL INFORMATION

3336™ F is a broad spectrum fungicide exhibiting preventative, curative, and systemic properties. It is useful on a wide variety of turf and ornamental disease problems. Apply 3336 F with ground or aerial equipment, using sufficient volume of spray to provide thorough coverage. Use the higher rates under conditions of severe disease pressure. Also, see local State Extension Service recommendations for application schedules. Chemigation instructions follow "Directions For Use." Read and follow these instructions carefully for this method of application.

Resistance Management: To avoid the development of tolerant strains of fungi, 3336 F should be used with fungicides of different modes of action. Cleary Chemical does not recommend the use of products containing benomyl or thiabendazole in combination or rotation with 3336 F. These utilize similar chemistry and mode of action and can contribute to development of disease tolerance. If, after using 3336 F as recommended, and the treatment is not effective, a tolerant strain of fungi may be present. Consult your local Cleary Chemical representative, your State Agricultural Experiment Station, or your State Cooperative Extension Service for proper disease identification and advice on the prompt use of some other suitable fungicide or disease control strategy. As long as recommended precautions are followed, 3336 F can remain useful for disease control.

Mixing Instructions: SHAKE WELL BEFORE USING. Some settling may occur during prolonged periods of non-use. High pH environments cause a shortened tank life for diluted product. The buffering of tank water to pH 6-7 prior to the addition of 3336 F is recommended. Add required amount of 3336 F to partially filled tank (1/2 total volume), agitate by mechanical or hydraulic means, add tank mix product if used (see below), agitate again and then add remaining required amount of water. Continuous agitation is recommended to keep the material in proper suspension. For best results, use spray mixture the same day it is prepared.

Tank Mixing Instructions: 3336 F is compatible with most commonly used pesticides. If tank mixing with other materials, add products in the following order: wettable powders, dry flowables, liquid flowables, emulsifiable concentrates, and soluble materials such as fertilizers. No claim of compatibility with other products is implied. Do not tank mix with copper-containing materials or with highly alkaline pesticides, such as Bordeaux mixture or lime sulfur. Consult the intended tank mix partner product label for appropriate application rates and use instructions. Follow the label directions for the most restrictive of label precautions and limitations. This product cannot be mixed with any product containing a label prohibition against such mixing. Read and observe the most restrictive precautionary statements and other information appearing on product labels used in mixtures. 3336 F may be applied in conjunction with chemically neutral liquid fertilizers. Application in conjunction with highly alkaline fertilizers, such as aqueous ammonia, may cause a degradation of the pesticide, resulting in reduced performance and should be avoided.

CHEMIGATION

GENERIC REQUIREMENTS

1. Apply this product only through the following types of systems: sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, solid set, or hand move; flood (basin); or drip (trickle) irrigation system(s). Do not apply this product through any other type of irrigation system.
2. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

3. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.
4. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
5. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

SPECIFIC REQUIREMENTS

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back towards the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being drawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

NON SPECIFIC REQUIREMENTS

1. Remove scale, pesticide residue, and other foreign matter from the chemical tank and entire injector system. Flush with clean water.
2. Prepare a suspension of product in the mix tank or stock bucket. Fill the tank with 1/2 or 3/4 of the desired amount of water. Start agitation and add the required amount of product to the solution along with the remaining volume of water.
3. Maintain a gentle agitation in the mix tank during application to assure a uniform suspension. Follow mixing instructions and tank mixing instructions previously indicated.

4. Start system and then uniformly inject the suspension of 3336™ F into the irrigation line so as to deliver the desired rate per acre. Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation system.
5. Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of a more dilute suspension per unit time.
6. The suspension of 3336 F should be injected with a positive displacement pump into the main line ahead of a right angle turn to insure adequate mixing.

SPRINKLER (OVERHEAD) CHEMIGATION

Observe all instructions in the General and Specific requirements sections above and the following additional requirements:

1. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Set sprinkler system to deliver 1/10 to 1/4 inches of water per acre. Volumes of water higher than this may reduce efficacy. Application of more than recommended quantities of irrigation water per acre may result in decreased product performance. Where sprinkler distribution patterns do not overlap sufficiently, unacceptable disease control may result.

When system connections or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product cannot be flushed and must be dismantled and drained in a center pivot system, block the nozzle set nearest the well pivot injection unit to prevent spray being applied to this area. Allow sufficient time for pesticides to be flushed through all lines and all nozzles before turning off irrigation water.

FLOOD (BASIN) CHEMIGATION

1. Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as a drop structure of weir box to decrease potential for water source contamination from backflow if water flow stops.
2. Systems utilizing a pressurized water and pesticide injection system must meet the following requirements: Observe all instructions in the General and Specific requirements sections above and items 1 and 2 of the sprinkler irrigation requirements.

DRIP (TRICKLE) CHEMIGATION

Observe all instructions in the General and Specific requirements sections above.

TURF APPLICATIONS

3336 F may be used on all fine turf applications such as commercial and residential lawns, parks, athletic fields, golf course greens, tees, and fairways, cemeteries, and sod farms of cool and warm season grasses such as Bentgrass, Bluegrass, Bermudagrass, Fescue, Ryegrass, St. Augustinegrass, Zoysiagrass, or their mixtures. 3336 F is not phytotoxic to any of the above mentioned grasses when used in accordance with the label. 3336 F is to be used for the prevention and control of the diseases mentioned below. It has both preventative and curative activity. Do not graze animals on treated turf. Do not feed clippings to livestock or poultry. Do not exceed a maximum of three (3) pounds active ingredient per 1,000 square feet per season from any thiophanate-methyl containing product.

Ground Application Instructions: Apply material with properly calibrated hand held, mechanical or motorized spray equipment or by chemigation through appropriate sprinkler irrigation systems. Spray uniformly over the area to be treated. Apply recommended amounts in sufficient water to obtain thorough coverage of treatment area (2-4 gal per 1,000 sq ft is suggested). When treating golf greens, always treat aprons and approaches. Use the highest recommended rate under conditions of severe disease pressure. For best results, apply after mowing or avoid mowing twelve hours after application. Lightly water the treatment area to move the fungicide into the active root zone with one to two tenths inch of water. Excessive irrigation may move application below active root zone and reduce application effectiveness. Green design and drainage will influence irrigation practices. When tank mixing with contact action fungicides for foliar diseases, applications should be allowed to dry on leaf surfaces. Normal watering may proceed after sprays have dried.

TURF APPLICATION

| Disease(s) Controlled | Rate of 3336 F fl oz/1,000 sq ft |
|--|-------------------------------------|
| Anthracnose, basal Colletotrichum graminicola | 4-8 |
| Anthracnose, foliar Colletotrichum graminicola | 2-4 |

For prevention in historic areas of disease pressure, apply twice at 14 day (d) intervals when soil temperature reaches 60°F. For curative control, apply when disease first appears and continue at 7-14 d intervals. Rotations and/or tank mix combinations with chlorothalonil or triadimefon can be utilized.

| | |
|---|-----|
| Bermudagrass Decline Gaeumannomyces graminis var. graminis | 4-8 |
| Take-All-Patch Gaeumannomyces graminis var. avenae | |

Apply in mid-July or when disease symptoms first appear and repeat at 7-14 d intervals for suppression. Use higher rates under most severe disease expression. Follow proper agronomic recommendations to maintain plant vigor.

| | |
|--|-----|
| Cool Season Brown Patch Rhizoctonia cerealis | 4-8 |
| Necrotic Ring Spot Leptosphaeria korrea | |
| Spring Dead Spot Leptosphaeria korrea | |

For prevention, apply in Fall before turf has stopped all growth activity. Apply second application in early Spring when soil temperatures reach 55-60°F or when disease first appears. For curative action, apply when disease first appears in early Spring and continue at 7-14 d intervals.

| | |
|--|-----|
| Coprinus Snow Mold Coprinus psychromorbidus | 4-8 |
|--|-----|

Apply 2 treatments at 21 d intervals in late Fall to early Winter, with the last application made just prior to first permanent snow cover. Rotations and/or tank mix combinations with Defend™ can be utilized.

| | |
|--------------------------------------|-----|
| Dollar Spot Moellerodiscus, Lanzia | 2-4 |
| Large Brown Patch Rhizoctonia solani | |
| Ascochyta Leaf Blight Ascochyta | |
| Copper Spot Gloeocercospora sorghi | |
| Fusarium Patch Fusarium nivale | |
| Red Thread Corticum fuciforme | |
| Zoysia Patch Rhizoctonia solani | |

Apply when disease first appears and continue at 7-14 d intervals. Rotations and/or tank mix combinations with chlorothalonil, iprodione, or ProtectT/O™ can be utilized.

| | |
|--|-----|
| Fusarium Blight Fusarium roseum, F. triticum | 4-8 |
|--|-----|

Apply when disease first appears 7-14 d intervals.

| | |
|---|-----|
| Gray Leaf Spot (Blast) Pyricularia grisea | 4-8 |
|---|-----|

Apply preventative application before expected period of disease development. Continue applications at 7-14 d intervals.

TURF APPLICATION (cont.)

| Disease(s) Controlled | Rate of 3336 F fl oz/1,000 sq ft |
|--|-------------------------------------|
| Leaf Spot Drechslera | 4-8 |
| Leaf, Crown, and Root Diseases Bipolaris, Curvularia, Exserohilum | |

Apply when disease first appears and make applications at 7-14 d intervals. Rotations and/or tank mix combinations with chlorothalonil, iprodione, or ProtectT/O are recommended under severe conditions.

| | |
|------------------------------------|-----|
| Pink Snow Mold Microdochium nivale | 2-4 |
|------------------------------------|-----|

Apply in late Fall to early Winter before turf has stopped all growth activity. A second application may be used in combination with chlorothalonil, Defend™, or Spotrete™ at recommended rates before snow cover or during Spring thaw.

| | |
|--------------------------|-----|
| Rusts Puccinia, Uromyces | 4-8 |
|--------------------------|-----|

Apply at 7-14 d intervals when disease first appears. Rotations and/or tank mix combinations with chlorothalonil or ProtectT/O are recommended.

| | |
|----------------------------------|-----|
| Stripe Smut Ustilago striiformis | 4-8 |
|----------------------------------|-----|

Apply at 7-14 d intervals when disease first appears. For prevention, apply in spring and fall.

| | |
|-------------------------------|-----|
| Summer Patch Magnaporthe poae | 4-8 |
|-------------------------------|-----|

For prevention, apply 3 applications starting late April or early May using 21 d intervals. Rotations and/or tank mix combinations may be used as part of the three application program. For suppression, apply at 7-14 d intervals when disease first appears.

HORTICULTURAL APPLICATIONS

NURSERY, GREENHOUSE, LANDSCAPE & INTERIORSCAPE

Annual and Perennial Flowers, Bedding Plants, Foliage Plants, Ground Covers, plus Deciduous and Evergreen Trees and Shrubs

3336 F is a broad spectrum systemic fungicide which controls a variety of foliar, stem, and root diseases on a wide range of commercially important plants. 3336 F is also effective as a pre-plant dip on cuttings and bulbs. For soil drench applications, best crop protection is achieved with preventative treatments repeated every 21-28 days. For foliar applications, begin treatments when disease first appears, or during suspected periods of disease incidence. Apply additional applications every 7-14 days or as otherwise instructed for the prevention or control of the listed diseases. Use of a wetting agent is recommended for plants that have leaves that are difficult to wet properly. Use of a spreader-sticker is recommended to enhance product performance in wet weather conditions or during periods of overhead irrigation. 3336 F may be used to control listed diseases on "backyard" (non-commercial) fruit trees such as almond, apple, apricot, cherry, nectarine, peach, pecan, plum, and prune trees. Do not apply within 1 day (24 hr) of harvest. 3336 F may be applied as a ground application using hand held, mechanical or motorized spray equipment, or as a chemigation spray or through an applicable sprinkler irrigation system; or as an aerial application where applicable. See specific instructions below. For foliar applications, do not exceed thirty-six (36) pounds active ingredient per acre per crop season from all thiophanate-methyl containing products. For soil drench applications, do not exceed 300 pounds active ingredient per acre per crop season from all thiophanate-methyl containing products.

Note: The "Directions For Use" of this product reflect the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied to any plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results. This product is not recommended for use on Swedish Ivy (*Neprolepis exaltata*), Boston Fern (*Plectranthus australis*), and Easter Cactus (*Hatiora gaertneri*).

Ground Application Instructions: Apply material with properly calibrated hand held, mechanical or motorized spray equipment or by chemigation thorough appropriate sprinkler irrigation, flood, or drip systems. Begin applications when disease first appears and repeat at 7-14 d intervals or as needed during the growing season. Use the shortest interval when conditions are unusually favorable for the development of disease. For hand held, mechanical, or motorized applications, mix 8-24 fl oz of 3336™ F per 100 gal water (0.5-1.5 teaspoons per gal) and apply as a full coverage spray to drip for the prevention and control of the diseases listed below. Spray volume may range up to 400 gallons of finished spray per acre depending upon plant species and plant growth stage. For applications through irrigation systems, refer to aerial use rates indicated in the foliar application chart. For small volume applications less than 100 gallons, divide recommended rate by 16 to get the number of teaspoons of 3336 F per gal.

SPECIAL INSTRUCTIONS FOR PROPORTIONAL INJECTORS

(e.g. Dosatron, Dosmatic, Anderson, and similar equipment)

Determine the treatment rate as indicated below in the ground application column for crop and pathogen. Determine the injection ratio for the individual system to be used for application. For systems using a 1:100 ratio, measure and add the exact amount of recommended material per 100 gallons to each gallon of water in a stock bucket or tank. For systems using a 1:200 ratio, multiply the recommended amount per 100 gallons by 2. For systems using a 1:50 ratio, divide the recommended amount per 100 gallons added by 2. For systems using a 1:16 ratio, divide the recommended amount per 100 gallons by 6. Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of a more dilute suspension per unit time. An injection ratio of 1:100 is recommended for most greenhouse and nursery systems.

Aerial Application Instructions: Apply with fixed wing or helicopter equipment to appropriate nursery sites (field and container). Use at rate indicated in sufficient water for thorough coverage or a minimum of 3 gallons per acre for low growing crops and a minimum of 10 gallons per acre for larger shrubs and trees. Use a spreader-sticker at recommended rates as needed to enhance product performance. Add product slowly to water in the spray tank with agitation or premix thoroughly in separate holding tank for concentrate or aircraft sprayers. Continuous agitation is required to keep the product in suspension.

FOLIAR APPLICATION

Use fluid ounce/acre rate for aerial applications and fluid ounce/100 gal rate for ground applications.

| Disease(s) Controlled | Rate of 3336 F oz. per | |
|--------------------------|------------------------|----------------|
| | 100 gallon Ground | acre Aerial |

Anthracnose 12-16 48-64
Colletotrichum

Apply as buds break or at first sign of disease. Repeat at 7-14 day (d) intervals as needed during disease period.

Black Spot of Rose 12-16 48-64
Diplocarpon rosae

Apply early summer or at first sign of disease. Repeat every 7-14 d as needed during disease period.

Brown Rot and Blight 12-16 48-64
Monilinia, Sclerotinia, Whetzellinia

Apply late Spring or at first sign of disease. Repeat every 7-14 d as needed during disease period.

Fusicladium and
Venturia Leaf Scabs on: 12-16 48-64
Crabapple, Hawthorn, Pear, Mountain Ash,
Pyracantha, etc.

Apply as buds break. Repeat every 7-14 d during disease period. Effective control requires coverage during leaf expansion. Rotations and/or tank mix combinations with ProtectT/O™ chlorothalonil or propiconazole can be utilized.

Leaf Spots and Blights 12-16 48-64
caused by: Ascochyta, Blumeriella, Botrytis, Cercospora,
Coccomyces, Corynespora, Curvularia, Didymellina,
Entomosporium, Fabraea, Fusarium, Ramularia,
Rhizoctonia, Marssonina, Mycosphaerella, Myrothecium,
Phoma, Physalospora, Schizothyrium, Septoria,
Sphaeloma

Apply when disease symptoms first appear. Repeat every 7-14 d during disease period. Rotations and/or tank mix combinations with ProtectT/O or chlorothalonil can be utilized.

Ovulinia Blight 8-16 32-64
Apply as flowers open. Repeat every 7-14 d during disease period.

Powdery Mildews 12-24 48-96
Erysiphe, Microsphaera, Phyllactinia, Podosphaera,
Oidium, Sphaerotheca

Apply when disease first appears and repeat every 7-14 d. Rotations and/or tank mix combinations with ProtectT/O or triadimefon can be utilized.

Rust Diseases caused by: 12-16 48-64
Puccinia, Gymnosporangium, Uromyces

Apply late Spring or when symptoms first appear. Repeat every 7-14 d during disease period. Rotations and/or tank mix combinations with ProtectT/O or chlorothalonil are recommended.

Tip Blight of Pine 16-24 64-96
Sphaeropsis sapinea, Diplodia pinea

Begin application in spring when new growth starts. Make a second application just before needles emerge from the sheath and a third application 14 d later. Thorough coverage is essential for optimal disease control.

| Disease(s) Controlled | Rate of 3336 F oz. per | |
|--------------------------|------------------------|----------------|
| | 100 gallon Ground | acre Aerial |

Twig Blights, Cankers, and Diebacks 16-24 64-96
Diaporthe, Kabatina, Phoma, Phomopsis

Apply when symptoms first appear. Repeat every 7-14 d during disease period.

SOIL DRENCH APPLICATION

| Disease(s) Controlled | Rate of 3336 F per | |
|--------------------------|--------------------|--|
| | 100 gallon | |

Stem, Crown, and Root Rots 8-16
caused by:

Botrytis, Cylindrocladium, Fusarium, Gliocladium,
Myrothecium, Penicillium, Ramularia, Rhizoctonia,
Sclerotinia

Black Root Rot Thielaviopsis

Apply as a drench or directed spray using hand held, mechanical or motorized spray equipment or as a chemigation drench or directed spray using applicable sprinkler irrigation systems after seeding or sticking of cuttings (8 fl oz) or after transplanting (12-16 fl oz) to propagation beds, containers, pots, trays, or nursery or landscape beds at a rate to thoroughly soak the growing media through the root zone. A general guide is 0.25-3.0 pints of finished mixture per sq ft depending on the media type and depth (about 4 fl oz per 4 inch pot or 8 fl oz per 6 inch pot). Repeat every 21-28 d for adequate crop protection.

Note: 3336 F does not control Pythium or Phytophthora. Tank mix combinations with metalaxyl, mefenoxam, etridiazole, propamocarb, or fosetyl-Al are required for the control of Pythium and Phytophthora.

PLANT DIP APPLICATION

| Disease(s) Controlled | Rate of 3336 F per | |
|--------------------------|--------------------|--|
| | 100 gallon | |

Plant or Cutting Diseases caused by: 16-24
Botrytis, Cylindrocladium, Fusarium, Gliocladium,
Myrothecium, Penicillium, Ramularia, Rhizoctonia,
Sclerotinia, Thielaviopsis

Immerse plants or cuttings for 10-15 min. Remove and allow to drain.

Note: Follow accepted hygiene practices to minimize the introduction and spread of water borne bacterial and water mold fungal diseases.

Bulb, Corm, and Rhizome Rots 16-24
caused by:

Botrytis, Cylindrocladium, Fusarium, Gliocladium,
Myrothecium, Penicillium, Ramularia, Rhizoctonia,
Sclerotinia, Thielaviopsis

Soak cleaned bulbs for 15-30 min in warm solution (80-85°F). For storage disease prevention, treat bulbs preferably within 48 h after digging. After treatment, dry well before storing. If bulbs are for forcing, treat bulbs that have been heat-cured.

Note: Follow accepted hygiene practices to minimize the introduction and spread of water borne bacterial and water mold fungal diseases.

LIMITED WARRANTY AND DISCLAIMER

CLEARY CHEMICAL CORPORATION warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the risks referred to therein. CLEARY CHEMICAL CORPORATION makes no other expressed or implied warranty of fitness or merchantability or any other expressed or implied warranty. In no case shall CLEARY CHEMICAL or seller be liable for consequential, special or indirect damages resulting from the use or handling of this product including, but not limited to, loss of profits, business reputation, or customers, labor costs, or other expenses incurred in planting or harvesting. CLEARY CHEMICAL and seller offer this product and the buyer and user accept it subject to the foregoing conditions of sale and warranty which may be varied only by agreement in writing signed by a duly authorized representative of CLEARY CHEMICAL CORPORATION.